



## Turbulence and Transport in Magnetically Confined Plasma

Juul Rasmussen, Jens; Garcia, O.E.; Madsen, Jens; Naulin, Volker; Nielsen, Anders Henry

*Publication date:*  
2012

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Juul Rasmussen, J., Garcia, O. E., Madsen, J., Naulin, V., & Nielsen, A. H. (2012). *Turbulence and Transport in Magnetically Confined Plasma*. Abstract from 6th International Conference on Solitons, Collapses and Turbulence, Akademgorodok, Novosibirsk, Russian Federation.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Turbulence and Transport in Magnetically Confined Plasma

**J. Juul Rasmussen<sup>1\*</sup>, O. E. Garcia<sup>2</sup>, J. Madsen<sup>1</sup>, V. Naulin<sup>1</sup>, A. H. Nielsen<sup>1</sup>**

<sup>1</sup>Association EURATOM-DTU, Technical University of Denmark, Department of Physics, DTU Risø Campus, P.O. Box 49, 4000 Roskilde, Denmark

<sup>2</sup>Department of Physics and Technology, University of Tromsø, 9037 Tromsø, Norway

\*e-mail address: jjra@fysik.dtu.dk

Transport mediated by turbulent is the dominating mechanism for plasma transport across a confining magnetic field in hot plasmas. The turbulent fluctuations arise spontaneously due to pressure gradients perpendicular to the magnetic field. Low frequency plasma turbulence is well described by the fluid model for plasma dynamics, and it is essentially quasi two-dimensional, i.e., the turbulence is strongly anisotropic with the main dynamics in the plane perpendicular to the magnetic field.

In toroidal magnetically confined plasmas the scrape-off layer, SOL, is the outer region of the plasma, where the magnetic field lines ends on material surfaces. It has the function of "scraping off" the plasma that is transported out through the edge region. The dynamics and transport in the edge region is essential for the global plasma dynamics, since all plasma has to leave through the edge. In addition, the character of the transport in the edge/SOL is essential for estimating the power deposition on the plasma facing components. It is well established from both experiments and models that the turbulence and transport in this region is inherently intermittent and involves large outbreak of hot plasma. These structures appear as filaments aligned with the magnetic field; they are often termed "blobs" and propagate far out in the SOL. The dynamics of these blobs, which resembles raising plumes in thermal convection, is well described by a two-dimensional convection model. The ensuing turbulent transport cannot be parameterized in terms of a simple diffusion coefficient (in the sense of Ficks law), but is characterized by strongly skewed Probability Density Functions (PDF) with broad (exponential) tails.

In this presentation results from numerical modelling of the transport in the Edge/SOL region of magnetically confined plasmas in toroidal devices will be reviewed and discussed in terms of experimental findings [1-4].

## References

- [1] O.E. Garcia, V. Naulin, A.H. Nielsen, J. Juul Rasmussen, "Computations of Intermittent Transport in Scrape-Off Layer Plasmas", *Phys. Rev. Lett.* **92**, 165003 (2004); "Turbulence and intermittent transport at the boundary of a magnetized plasmas", *Phys. Plasmas* **12**, 090701 (2005).
- [2] O.E. Garcia, J. Horacek, R.A. Pitts, A.H. Nielsen, W. Fundamenski, J.P. Graves, V. Naulin, J. Juul Rasmussen, "Interchange turbulence in the TCV scrape-off layer", *Plasma Phys. Control. Fusion* **48**, L1 (2006).
- [3] O.E. Garcia, J. Horacek, R.A. Pitts, A.H. Nielsen, W. Fundamenski, V. Naulin, J. Juul Rasmussen, "Fluctuations and transport in the TCV scrape-off layer", *Nucl. Fusion* **47**, 667 (2007).
- [4] J. Madsen, O.E. Garcia, J.S. Larsen, V. Naulin, A.H. Nielsen, J. Juul Rasmussen, "The influence of finite Larmor radius effects on the radial interchange motions of plasma filaments", *Phys. Plasmas* **18**, 112504 (2011).